

Gauge-invariant description of several (2+1)-dimensional integrable nonlinear evolution equations

V. G. Dubrovsky and A. V. Gramolin

Novosibirsk State Technical University, Novosibirsk, Russia

E-mail: dubrovsky@academ.org and gramolin@gmail.com

Abstract

We obtain new gauge-invariant forms of two-dimensional integrable systems of nonlinear equations: the Sawada–Kotera and Kaup–Kuperschmidt system, the generalized system of dispersive long waves, and the Nizhnik–Veselov–Novikov system. We show how these forms imply both new and well-known two-dimensional integrable nonlinear equations: the Sawada–Kotera equation, Kaup–Kuperschmidt equation, dispersive long-wave system, Nizhnik–Veselov–Novikov equation, and modified Nizhnik–Veselov–Novikov equation. We consider Miura-type transformations between nonlinear equations in different gauges.

Keywords: Sawada–Kotera equation, Kaup–Kuperschmidt equation, generalized dispersive long-wave equation, Davey–Stewartson equation, Nizhnik–Veselov–Novikov equation

1 Introduction

The fundamental methods based on gauge transformations and the concept of gauge invariance are currently widely used in physics and mathematics, in particular, in the theory of integrable nonlinear equations. The first applications of these methods in the theory of integrable nonlinear equations were proposed in [1–7] (also see [8–13] and the references therein).

A great many gauge-equivalent pairs of integrable models have been found. In the one-dimensional case, these include the nonlinear Schrödinger equation and Heisenberg ferromagnet equation, the equations describing the massive Thirring model and the two-dimensional relativistic field theory, and the Korteweg–de Vries (KdV) and modified KdV (mKdV) equations. In the two-dimensional case, the currently most widely known gauge-equivalent pairs of equations are the Kadomtsev–Petviashvili (KP) and modified KP (mKP) equations and the Davey–Stewartson and Ishimori equations (see [8–14] and the references therein).

We introduce the required terms and illustrate the unifying role of gauge transformations and gauge invariance with the well-known example of the interaction between a charged spinless particle and an external electromagnetic field with the vector and scalar potentials $\vec{A}(\vec{r}, t)$ and $\psi(\vec{r}, t)$,

$$i\hbar\psi_t = \frac{(\hat{\vec{p}} - q\vec{A})^2}{2m}\psi + q\phi\psi, \quad (1.1)$$

where $\widehat{\vec{p}} = -i\hbar\vec{\nabla}$ is the particle momentum operator. The particle coupling to external fields has the well-known gauge-invariant form.

From the standpoint of the inverse scattering method, Eq. (1.1) is an auxiliary linear problem: a linear partial differential equation with variable coefficients for the wave function ψ . Under the gauge transformation

$$\psi \rightarrow \psi' = g^{-1}\psi, \quad \psi = g\psi' = \exp\left[\frac{i\chi(\vec{r}, t)q}{\hbar}\right]\psi', \quad (1.2)$$

Eq. (1.1) preserves its form if the potentials $\vec{A}$ and ϕ are transformed as

$$\vec{A} \rightarrow \vec{A}' = \vec{A} - \vec{\nabla}\chi, \quad \phi \rightarrow \phi' = \phi + \chi_t. \quad (1.3)$$

Eliminating the gauge function χ from (1.3), we obtain the relations

$$[\vec{\nabla} \times \vec{A}'] = [\vec{\nabla} \times \vec{A}], \quad -\vec{\nabla}\phi' - \frac{\partial \vec{A}'}{\partial t} = -\vec{\nabla}\phi - \frac{\partial \vec{A}}{\partial t}, \quad (1.4)$$

which mean that the quantities

$$\vec{B} \stackrel{\text{def}}{=} [\vec{\nabla} \times \vec{A}], \quad \vec{E} \stackrel{\text{def}}{=} -\vec{\nabla}\phi - \frac{\partial \vec{A}}{\partial t}, \quad (1.5)$$

which are well known in electrodynamics as the magnetic field induction $\vec{B}$ and electric field strength $\vec{E}$, are invariants of gauge transformations. Moreover, definitions (1.5) for the invariants $\vec{B}$ and $\vec{E}$ imply the subsystem of equations

$$\text{div}\vec{B} = \vec{\nabla} \cdot \vec{B} = 0, \quad \vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}, \quad (1.6)$$

which is a (sourceless) subsystem of the fundamental system of Maxwell equations. Thus, starting from the principle of local gauge invariance, we can obtain the gauge-invariant non-stationary Schrödinger equation for a charged spinless particle in an external electromagnetic field, the gauge invariants $\vec{B}$ and $\vec{E}$, and a (sourceless) subsystem of fundamental Maxwell equations. Similar considerations are applicable to the case of integrable nonlinear equations.

2 Gauge-invariant integrable system of KP and mKP equations

We illustrate the methods based on the notion of gauge invariance with the well-known example of an integrable system of nonlinear KP and mKP equations. Auxiliary linear problems for the KP and mKP equations are particular cases of the following linear problems (with the respective gauges $C(u_0 \neq 0, u_1 = 0)$ and $C(u_0 = 0, u_1 \neq 0)$ for the KP and mKP equations):

$$L_1\psi = (\sigma\partial_y + \partial_x^2 + u_1\partial_x + u_0)\psi = 0, \quad (2.1)$$

$$L_2\psi = (\partial_t + 4\partial_x^3 + v_2\partial_x^2 + v_1\partial_x + v_0)\psi = 0, \quad (2.2)$$

where the constant values $\sigma = i$ or $\sigma^2 = -1$ and $\sigma = 1$ or $\sigma^2 = 1$ respectively correspond to the cases of KP-I or mKP-I and KP-II or mKP-II equations. Because of (2.1) and (2.2), the compatibility condition in the Lax form $[L_1, L_2] = 0$ leads to a system of evolution equations for the field variables u_1 and u_0 [5, 7],

$$\begin{aligned} u_{1t} + u_{1xxx} - \frac{3}{2}u_1^2u_{1x} - 3\sigma u_{1x}\partial_x^{-1}u_{1y} + 3\sigma^2\partial_x^{-1}u_{1yy} + \\ + 6u_{0xx} - 6\sigma u_{0y} + 6(u_0u_1)_x - 2v_{0x} = 0, \\ u_{0t} + 4\sigma u_{0xxx} + 6u_0u_{0x} + 6u_1u_{0xx} + \frac{3}{2}u_1^2u_{0x} + 3u_{0x}u_{1x} - \\ - 3\sigma u_{0x}\partial_x^{-1}u_{1y} - u_1v_{0x} - v_{0xx} - \sigma v_{0y} = 0. \end{aligned} \quad (2.3)$$

For the gauge $C(u_0 \neq 0, u_1 = 0)$ with the choice of the variable $v_0 = 3u_{0x} - 3\sigma\partial_x^{-1}u_{0y}$, system (2.3) implies the KP equation [1, 15]

$$u_{0t} + u_{0xxx} + 6u_0u_{0x} + 3\sigma^2\partial_x^{-1}u_{0yy} = 0. \quad (2.4)$$

For the gauge $C(u_0 = 0, u_1 \neq 0)$ with the choice of the variable $v_0 = \text{const}$, system (2.3) implies the mKP equation

$$u_{1t} + u_{1xxx} - \frac{3}{2}u_1^2u_{1x} - 3\sigma u_{1x}\partial_x^{-1}u_{1y} + 3\sigma^2\partial_x^{-1}u_{1yy} = 0. \quad (2.5)$$

System of equations (2.3) is called the system of KP-mKP equations.

In terms of the pure gauge variable ρ and the invariant w_0 of the gauge transformations $\psi \rightarrow \psi' = g^{-1}\psi$ determined by the expressions

$$u_1(x, y, t) = 2\frac{\rho_x}{\rho}, \quad w_0 = u_0 - \frac{1}{2}u_{1x} - \frac{1}{4}u_1^2 - \frac{\sigma}{2}\partial_x^{-1}u_{1y}, \quad (2.6)$$

system of equations (2.3) becomes

$$\rho_t + 4\rho_{xxx} + 6\rho_x w_0 + 3\rho w_{0x} - 3\sigma\partial_x^{-1}w_{0y} - \rho v_0 = 0, \quad (2.7)$$

$$w_{0t} + w_{0xxx} + 6w_0w_{0x} + 3\sigma^2\partial_x^{-1}w_{0yy} = 0. \quad (2.8)$$

System (2.7), (2.8) is the manifestly gauge-invariant form of system of KP-mKP equations (2.3). This system has the following structure. It contains gauge-invariant KP equation (2.8) for the gauge invariant w_0 and Eq. (2.7) for the pure gauge variable ρ with additional terms containing the gauge invariant w_0 and the additional field variable v_0 . If the invariant is zero, $w_0 = 0$, then KP-mKP system (2.7), (2.8) reduces to the linear evolution equation for the pure gauge variable ρ :

$$\rho_t + 4\rho_{xxx} - \rho v_0 = 0. \quad (2.9)$$

We use the gauge invariant w_0 to obtain the Miura-type transformation from Eq. (2.8),

$$w_0 = u_0 = -\frac{1}{2}u_{1x} - \frac{1}{4}u_1^2 - \frac{\sigma}{2}\partial_x^{-1}u_{1y}, \quad (2.10)$$

which relates the solutions u_0 (for the gauge $C(u_0 \neq 0, u_1 = 0)$) and u_1 (for the gauge $C(u_0 = 0, u_1 \neq 0)$) of KP equations (2.4) and mKP equations (2.5). This also follows from Eq. (2.8) written for the gauges $C(u_0 \neq 0, u_1 = 0)$ and $C(u_0 = 0, u_1 \neq 0)$:

$$u_{0t} + u_{0xxx} + 6u_0u_{0x} + 3\sigma^2\partial_x^{-1}u_{0yy} = -\frac{1}{2}\left(\partial_x + \frac{1}{2}u_1 + \sigma\partial_x^{-1}\partial_y\right)\left(u_{1t} + u_{1xxx} - \frac{3}{2}u_1^2u_{1x} - 3\sigma u_{1x}\partial_x^{-1}u_{1y} + 3\sigma^2\partial_x^{-1}u_{1yy}\right) = 0. \quad (2.11)$$

We stress that the separation of the physical and the pure gauge degrees of freedom in integrable nonlinear equations and their gauge-invariant formulation can be used to study the structure of these equations and the relations between their different gauge-equivalent realizations.

3 Manifestly gauge-invariant integrable system of two-dimensional Kaup–Kuperschmidt and Sawada–Kotera equations

In this section, we briefly discuss the results obtained up to now concerning the manifestly gauge-invariant formulation of two-dimensional integrable generalizations of the Kaup–Kuperschmidt (2DKK) and Sawada–Kotera (2DSK) nonlinear equations. The auxiliary linear problems for these equations are particular cases (in the gauge $C(u_2 = 0, u_1 \neq 0, u_0 = u_{1x}/2)$ for the 2DKK equation and in the gauge $C(u_2 = 0, u_1 \neq 0, u_0 = 0)$ for the 2DSK equation) of the problems [7]

$$\begin{aligned} L_1\psi &= (\sigma\partial_y + \partial_x^3 + u_2\partial_x^2 + u_1\partial_x + u_0)\psi = 0, \\ L_2\psi &= (\partial_t + \kappa\partial_x^5 + v_4\partial_x^4 + v_3\partial_x^3 + v_2\partial_x^2 + v_1\partial_x + v_0)\psi = 0. \end{aligned} \quad (3.1)$$

In terms of the pure gauge variable ρ and the invariants w_0 and w_1 of the gauge transformations $\psi \rightarrow \psi' = g^{-1}\psi$ given by the expressions

$$u_2 = 3\frac{\rho_x}{\rho}, \quad w_1 = u_1 - u_{2x} - \frac{1}{3}u_2^2, \quad (3.2)$$

$$w_0 = u_0 - \frac{1}{3}u_1u_2 - \frac{1}{3}u_{2xx} + \frac{2}{27}u_2^3 - \frac{\sigma}{3}\partial_x^{-1}u_{2y}, \quad (3.3)$$

we obtain the manifestly gauge-invariant integrable system of 2DKK–2DSK nonlinear equations

$$\begin{aligned} \rho_t + \kappa\rho_{xxxxx} - \rho v_0 + \frac{5}{3}\kappa(\rho_{xx}w_1)_x + \frac{5}{3}\kappa(\rho_xw_0)_x + \frac{5}{9}\kappa\rho_xw_1^2 + \frac{10}{9}\kappa\rho_xw_{1xx} + \\ + \frac{10}{9}\kappa\rho w_{0xx} + \frac{10}{9}\kappa\rho w_0w_1 - \frac{5}{9}\kappa\sigma\rho_x\partial_x^{-1}w_{1y} - \frac{5}{9}\kappa\sigma\rho\partial_x^{-1}w_{0y} = 0, \end{aligned} \quad (3.4)$$

$$\begin{aligned} w_{1t} - \frac{1}{9}\kappa w_{1xxxxx} - \frac{5}{9}\kappa(w_1w_{1xx})_x - \frac{5}{3}\kappa(w_0w_{1x})_x - \frac{5}{9}\kappa w_1^2w_{1x} + \frac{10}{3}\kappa w_0w_{0x} - \\ - \frac{5}{9}\kappa\sigma w_{1xxy} - \frac{5}{9}\kappa\sigma w_1w_{1y} + \frac{5}{9}\kappa\sigma^2\partial_x^{-1}w_{1yy} - \frac{5}{9}\kappa\sigma w_{1x}\partial_x^{-1}w_{1y} = 0, \end{aligned} \quad (3.5)$$

$$\begin{aligned} w_{0t} - \frac{1}{9}\kappa w_{0xxxxx} - \frac{5}{9}\kappa(w_0w_1)_{xxx} - \frac{5}{9}\kappa(w_0w_{1xx})_x + \frac{5}{3}\kappa(w_0w_{0x})_x - \frac{5}{9}\kappa(w_0w_1^2)_x - \\ - \frac{5}{9}\kappa\sigma w_{0xxy} - \frac{10}{9}\kappa\sigma w_0w_{1y} - \frac{5}{9}\kappa\sigma w_1w_{0y} + \frac{5}{9}\kappa\sigma^2\partial_x^{-1}w_{0yy} - \frac{5}{9}\kappa\sigma w_{0x}\partial_x^{-1}w_{1y} = 0. \end{aligned} \quad (3.6)$$

This system consists of gauge-invariant system of equations (3.5), (3.6) for the gauge invariants w_0 and w_1 and Eq. (3.4) for the pure gauge variable ρ with additional terms containing the gauge invariants w_0 and w_1 and the additional variable v_0 . For the zero-valued invariants $w_0 = 0$ and $w_1 = 0$, the *2DKK–2DSK* system given by (3.5) and (3.6) reduces to the linear evolution equation for the pure gauge variable ρ :

$$\rho_t + \kappa \rho_{xxxxx} - \rho v_0 = 0. \quad (3.7)$$

Gauge-invariant system of equations (3.5), (3.6) for the invariants w_0 and w_1 coincides in form with the system obtained in [7]. This system admits the following reductions:

1. In the case $w_0 = 0$, we obtain an equation for the invariant w_1 exactly coinciding with the *2DSK* equation [7, 16].
2. In the case $w_0 = w_{1x}/2$, we obtain an equation for the invariant w_1 exactly coinciding with the *2DKK* equation [7, 16].

Obviously, the *2DKK* and *2DSK* equations, as equations belonging to different sets of invariants, are not gauge-equivalent to each other.

4 A manifestly gauge-invariant integrable two-dimensional generalized dispersive long-wave system

The gauge-invariant formulation of integrable systems of nonlinear equations can be obtained in all cases where the gauge freedom is taken into account in the corresponding auxiliary linear problems. In [17], such a formulation was obtained for the auxiliary linear problems

$$L_1 \psi = (\partial_{\xi\eta}^2 + u_1 \partial_\xi + v_1 \partial_\eta + u_0) \psi = 0, \quad (4.1)$$

$$L_2 \psi = (\partial_t + \kappa_1 \partial_\xi^2 + \kappa_2 \partial_\eta^2 + \tilde{u}_1 \partial_\xi + \tilde{v}_1 \partial_\eta + v_0) \psi = 0, \quad (4.2)$$

where κ_1 and κ_2 are constants, $\xi = x + \sigma y$ and $\eta = x - \sigma y$ are spatial variables, $\sigma^2 = \pm 1$, and the derivatives are $\partial_\xi = \partial/\partial\xi$, $\partial_\eta = \partial/\partial\eta$, $\partial_\xi^2 = \partial^2/\partial\xi^2$, etc. Such a choice of auxiliary linear problems (4.1) and (4.2) leads to the well-known two-dimensional generalized dispersive long-wave (*2DgDLW*) equation [18], the system of Davey–Stewartson (DS) equations [19], their reductions, and some other equations. All the listed well-known integrable nonlinear equations were previously obtained from the compatibility condition for auxiliary linear problems (4.1) and (4.2) in the form of the Manakov triad representation [20]

$$[L_1, L_2] = B L_1. \quad (4.3)$$

To obtain the manifestly gauge-invariant formulation of the corresponding integrable system of nonlinear equations, it is convenient to use the classical gauge invariants w_2 , $\tilde{w}_2$, and w_1 ,

$$\begin{aligned} w_2 &\stackrel{\text{def}}{=} u_0 - u_{1\xi} - u_1 v_1 = u'_0 - u'_{1\xi} - u'_1 v'_1, \\ \tilde{w}_2 &\stackrel{\text{def}}{=} u_0 - v_{1\eta} - u_1 v_1 = u'_0 - v'_{1\eta} - u'_1 v'_1, \\ w_1 &\stackrel{\text{def}}{=} u_{1\xi} - v_{1\eta} = u'_{1\xi} - v'_{1\eta}, \end{aligned} \quad (4.4)$$

and the pure gauge variable ρ related to the field variable $u_1(\xi, \eta, t)$ as

$$u_1 \stackrel{\text{def}}{=} (\log \rho)_\eta. \quad (4.5)$$

The variable ρ corresponds to the pure gauge degrees of freedom and is transformed according to the simple law $\rho \rightarrow \rho' = g\rho$ under the gauge transformations $\psi \rightarrow \psi' = g^{-1}\psi$.

We note that the invariants w_2 and $\tilde{w}_2$ in auxiliary problem (4.1) are the Laplace invariants $h = w_2$ and $k = \tilde{w}_2$ for the corresponding classical differential equation (see, e.g., the well-known Forsyth monograph on differential equations [21]).

In the case under study with auxiliary linear problem (4.2) determining the time evolution, the corresponding integrable system of nonlinear equations in terms of the variables ρ , w_1 , and w_2 becomes

$$\rho_t = -\kappa_1 \rho_{\xi\xi} - \kappa_2 \rho_{\eta\eta} - 2\kappa_1 \rho \partial_\eta^{-1} w_{2\xi} + 2\kappa_2 \rho_\eta \partial_\xi^{-1} w_1 + v_0 \rho, \quad (4.6)$$

$$w_{1t} = -\kappa_1 w_{1\xi\xi} + \kappa_2 w_{1\eta\eta} - 2\kappa_1 w_{2\xi\xi} + 2\kappa_2 w_{2\eta\eta} - 2\kappa_1 (w_1 \partial_\eta^{-1} w_1)_\xi + 2\kappa_2 (w_1 \partial_\xi^{-1} w_1)_\eta, \quad (4.7)$$

$$w_{2t} = \kappa_1 w_{2\xi\xi} - \kappa_2 w_{2\eta\eta} - 2\kappa_1 (w_2 \partial_\eta^{-1} w_1)_\xi + 2\kappa_2 (w_2 \partial_\xi^{-1} w_1)_\eta. \quad (4.8)$$

Gauge-invariant subsystem (4.7), (4.8) for the invariants $w_1 = u_{1\xi} - v_{1\eta}$ and $w_2 = u_0 - u_{1\xi} - u_1 v_1$ of system of equations (4.6)–(4.8) with $u_1 = 0$, $v_1 = -q/2$, and $u_0 = (1 + r - q_\eta)/4$ in terms of the variables q and r becomes

$$\begin{aligned} q_t &= -\kappa_1 \partial_\eta^{-1} r_{\xi\xi} + \kappa_2 r_\eta - \frac{\kappa_1}{2} (q^2)_\xi + \kappa_2 q_\eta \partial_\xi^{-1} q_\eta, \\ r_t &= -\kappa_1 q_\xi + \kappa_2 \partial_\xi^{-1} q_{\eta\eta} - \kappa_1 q_{\eta\xi\xi} + \kappa_2 q_{\eta\eta\eta} - \kappa_1 (rq)_\xi + \kappa_2 (r \partial_\xi^{-1} q_\eta)_\eta. \end{aligned} \quad (4.9)$$

In the particular case where $\kappa_2 = 0$, system of equations (4.9) reduces to the well-known integrable $2DgDLW$ system [18]

$$\begin{aligned} q_{t\eta} &= -\kappa_1 r_{\xi\xi} - \frac{\kappa_1}{2} (q^2)_{\xi\eta}, \\ r_{t\xi} &= -\kappa_1 (qr + q + q_{\xi\eta})_{\xi\xi}. \end{aligned} \quad (4.10)$$

In the one-dimensional limit $\xi = \eta$, system (4.9) ($\kappa_1 - \kappa_2 = 1$) and system (4.10) ($\kappa_1 = 1$) reduce to the well-known dispersive long-wave equation (see, e.g., [22]). Therefore, integrable system of nonlinear equations (4.6)–(4.8) can be called the $2DgDLW$ system.

Integrable system (4.6)–(4.8) in terms of the variables $\phi = \log \rho$, w_1 , and w_2 becomes

$$\phi_t = -\kappa_1 \phi_{\xi\xi} - \kappa_2 \phi_{\eta\eta} - \kappa_1 (\phi_\xi)^2 - \kappa_2 (\phi_\eta)^2 - 2\kappa_1 \partial_\eta^{-1} w_{2\xi} + 2\kappa_2 \phi_\eta \partial_\xi^{-1} w_1 + v_0, \quad (4.11)$$

$$w_{1t} = -\kappa_1 w_{1\xi\xi} - \kappa_2 w_{1\eta\eta} - 2\kappa_1 w_{2\xi\xi} + 2\kappa_2 w_{2\eta\eta} - 2\kappa_1 (w_1 \partial_\eta^{-1} w_1)_\xi + 2\kappa_2 (w_1 \partial_\xi^{-1} w_1)_\eta, \quad (4.12)$$

$$w_{2t} = \kappa_1 w_{2\xi\xi} - \kappa_2 w_{2\eta\eta} - 2\kappa_1 (w_2 \partial_\eta^{-1} w_1)_\xi + 2\kappa_2 (w_2 \partial_\xi^{-1} w_1)_\eta. \quad (4.13)$$

Integrable system (4.6)–(4.8) in terms of the variables $\phi = \log \rho$, w_2 , and $\tilde{w}_2 = w_2 + w_1$ becomes more symmetric,

$$\phi_t = -\kappa_1 \phi_{\xi\xi} - \kappa_2 \phi_{\eta\eta} - \kappa_1 (\phi_\xi)^2 - \kappa_2 (\phi_\eta)^2 - 2\kappa_1 \partial_\eta^{-1} w_{2\xi} + 2\kappa_2 \phi_\eta \partial_\xi^{-1} w_1 + v_0, \quad (4.14)$$

$$w_{2t} = \kappa_1 w_{2\xi\xi} - \kappa_2 w_{2\eta\eta} - 2\kappa_1 (w_2 \partial_\eta^{-1} (\tilde{w}_2 - w_2))_\xi + 2\kappa_2 (w_2 \partial_\xi^{-1} (\tilde{w}_2 - w_2))_\eta, \quad (4.15)$$

$$\tilde{w}_{2t} = -\kappa_1 \tilde{w}_{2\xi\xi} + \kappa_2 \tilde{w}_{2\eta\eta} - 2\kappa_1 (\tilde{w}_2 \partial_\eta^{-1} (\tilde{w}_2 - w_2))_\xi + 2\kappa_2 (\tilde{w}_2 \partial_\xi^{-1} (\tilde{w}_2 - w_2))_\eta. \quad (4.16)$$

All the mutually equivalent integrable systems of $2DgDLW$ nonlinear equations considered above, (4.6)–(4.8), (4.11)–(4.13), and (4.14)–(4.16) have the following common characteristic gauge structure:

- a. They contain manifestly gauge-invariant subsystems (4.7), (4.8) and (4.12), (4.13) of nonlinear equations for the gauge invariants w_1 and w_2 (or, equivalently, subsystem (4.15), (4.16) for the gauge invariants w_2 and $\tilde{w}_2$).
- b. They contain Eq. (4.6) for the pure gauge variable ρ (or Eq. (4.11) for the variable $\phi = \log \rho$), which satisfies a simple transformation law $\rho \rightarrow \rho' = g\rho$ with additional terms containing the gauge invariants and the additional field variable v_0 .

Such a structure of the 2DgDLW system reflects a significant gauge freedom in auxiliary linear problems (4.1) and (4.2).

We consider several particular gauges of systems of 2DgDLW equations (4.6)–(4.8), (4.11)–(4.13), and (4.14)–(4.16). We let $C(u_1, v_1, u_0)$ denote the gauge in general position. In the gauge $C(u_1 = \phi_\eta, v_1 = \phi_\xi, u_0 = \phi_{\xi\eta} + \phi_\xi\phi_\eta)$, which by definitions (4.4) of the invariants corresponds to the zero values of the invariants w_1 and w_2 ,

$$w_1 = u_{1\xi} - v_{1\eta} = 0, \quad w_2 = u_0 - u_{1\xi} - u_1v_1 = 0, \quad \tilde{w}_2 = 0, \quad (4.17)$$

system of 2DgDLW equations (4.14)–(4.16) reduces to the two-dimensional Burgers equation in potential form

$$\phi_t = -\kappa_1\phi_{\xi\xi} - \kappa_2\phi_{\eta\eta} - \kappa_1(\phi_\xi)^2 - \kappa_2(\phi_\eta)^2 + v_0, \quad (4.18)$$

or, in terms of the variable ρ related to the Hopf–Cole transformation $\phi = \log \rho$, to the linear diffusion equation

$$\rho_t = -\kappa_1\rho_{\xi\xi} - \kappa_2\rho_{\eta\eta} + v_0\rho. \quad (4.19)$$

It follows from the above construction that Eq. (4.18) (or Eq. (4.19)) is the compatibility condition for auxiliary problems (4.1) and (4.2) in Lax form,

$$[L_1, L_2] = B(w_1)L_1 \equiv 0. \quad (4.20)$$

In another simple gauge, $C(u_1 = \phi_\eta, v_1 = 0, u_0 = 0)$, which by definitions (4.4) corresponds to the invariants

$$w_1 = \phi_{\xi\eta}, \quad w_2 = -\phi_{\xi\eta}, \quad \tilde{w}_2 = 0, \quad (4.21)$$

system of 2DgDLW equations (4.14)–(4.16) again reduces to the single Burgers-type equation in potential form

$$\phi_t = \kappa_1\phi_{\xi\xi} - \kappa_2\phi_{\eta\eta} - \kappa_1(\phi_\xi)^2 + \kappa_2(\phi_\eta)^2 + v_0. \quad (4.22)$$

This equation can be linearized by the Hopf–Cole transformation $\phi = -\log \rho$ to the corresponding linear equation

$$\rho_t = \kappa_1\rho_{\xi\xi} - \kappa_2\rho_{\eta\eta} - \rho v_0. \quad (4.23)$$

In the gauge $C(u_1 = 0, v_1 = -q_\xi/q, u_0 = pq)$, it follows from (4.4) that the invariants w_1 , w_2 , and $\tilde{w}_2$ are given by the expressions

$$w_1 = (\log q)_{\xi\eta}, \quad w_2 = u_0 = pq, \quad \tilde{w}_2 = pq + (\log q)_{\xi\eta}, \quad (4.24)$$

the variable ρ according to (4.5) has a constant value, and the variable ϕ is equal to zero. According to (4.14), for the variable v_0 , we have

$$v_0 = -2\kappa_1\partial_\eta^{-1}w_{2\xi} = -2\kappa_1\partial_\eta^{-1}(pq)_\xi. \quad (4.25)$$

After some calculations in the case under study, system of 2DgDLW equations (4.14)–(4.16) implies the well-known system of DS equations [19] for the variables p and q :

$$\begin{aligned} p_t &= \kappa_1 p_{\xi\xi} - \kappa_2 p_{\eta\eta} + 2\kappa_1 p \partial_\eta^{-1}(pq)_\xi - 2\kappa_2 p \partial_\xi^{-1}(pq)_\eta, \\ q_t &= -\kappa_1 q_{\xi\xi} + \kappa_2 q_{\eta\eta} - 2\kappa_1 q \partial_\eta^{-1}(pq)_\xi + 2\kappa_2 q \partial_\xi^{-1}(pq)_\eta. \end{aligned} \quad (4.26)$$

In the gauge $C(u_1 = p_\eta, v_1 = q_\xi, u_0 = p_\eta q_\xi)$, it follows from (4.4) that the invariants can be expressed in terms of the variables q and p as

$$w_1 = p_{\xi\eta} - q_{\xi\eta}, \quad w_2 = -p_{\xi\eta}, \quad \tilde{w}_2 = -q_{\xi\eta}. \quad (4.27)$$

We substitute w_1 , w_2 , and $\tilde{w}_2$ given by (4.27) in system (4.14)–(4.16) and obtain three equations for the variables p and q . Equation (4.14) for $\phi \equiv p$ implies the first equation

$$p_t = \kappa_1 p_{\xi\xi} - \kappa_2 p_{\eta\eta} - \kappa_1 (p_\xi)^2 + \kappa_2 (p_\eta)^2 - 2\kappa_2 p_\eta q_\eta + v_0. \quad (4.28)$$

The other two equations follow from Eqs. (4.15) and (4.16) for the invariants w_2 and $\tilde{w}_2$ and can be expressed in terms of the variables p and q as

$$p_t = \kappa_1 p_{\xi\xi} - \kappa_2 p_{\eta\eta} - \kappa_1 (p_\xi)^2 + \kappa_2 (p_\eta)^2 + 2\kappa_1 \partial_\eta^{-1}(p_{\xi\eta} q_\xi) - 2\kappa_2 \partial_\xi^{-1}(p_{\xi\eta} q_\eta), \quad (4.29)$$

$$q_t = -\kappa_1 q_{\xi\xi} + \kappa_2 q_{\eta\eta} + \kappa_1 (q_\xi)^2 - \kappa_2 (q_\eta)^2 - 2\kappa_1 \partial_\eta^{-1}(q_{\xi\eta} p_\xi) + 2\kappa_2 \partial_\xi^{-1}(q_{\xi\eta} p_\eta). \quad (4.30)$$

Equations (4.28) and (4.29) are compatible under the choice of the variable

$$v_0 = 2\kappa_1 \partial_\eta^{-1}(p_{\xi\eta} q_\xi) + 2\kappa_2 \partial_\xi^{-1}(q_{\xi\eta} p_\eta). \quad (4.31)$$

In this case, system of three equations (4.28)–(4.30) reduces to system of two equations (4.29) and (4.30), which contains the derivatives $p_{\xi\eta} q_\xi$, $p_{\xi\eta} q_\eta$, etc., in nonlocal terms.

Similarly, in the gauge $C(u_1 = p_\eta, v_1 = q_\xi, u_0 = 0)$, it follows from (4.4) that the invariants w_1 , w_2 , and $\tilde{w}_2$ can be expressed as

$$w_1 = p_{\xi\eta} - q_{\xi\eta}, \quad w_2 = -p_{\xi\eta} - p_\eta q_\xi, \quad \tilde{w}_2 = -q_{\xi\eta} - p_\eta q_\xi. \quad (4.32)$$

Equation (4.11) for $\phi \equiv p$ with (4.32) taken into account becomes

$$p_t = \kappa_1 p_{\xi\xi} - \kappa_2 p_{\eta\eta} - \kappa_1 (p_\xi)^2 + \kappa_2 (p_\eta)^2 - 2\kappa_2 p_\eta q_\eta + 2\kappa_1 \partial_\eta^{-1}(p_\eta q_\xi)_\xi + v_0. \quad (4.33)$$

Substituting expressions (4.32) for w_1 and w_2 , we transform Eq. (4.12) as

$$\begin{aligned} p_t - q_t &= \kappa_1 (p + q)_{\xi\xi} - \kappa_2 (p + q)_{\eta\eta} - \kappa_1 (p_\xi - q_\xi)^2 + \\ &+ \kappa_2 (p_\eta - q_\eta)^2 + 2\kappa_1 \partial_\eta^{-1}(p_\eta q_\xi)_\xi - 2\kappa_2 \partial_\xi^{-1}(p_\eta q_\xi)_\eta. \end{aligned} \quad (4.34)$$

Subtracting Eq. (4.34) from (4.33), we obtain the evolution equation for the variable q :

$$q_t = -\kappa_1 q_{\xi\xi} + \kappa_2 q_{\eta\eta} + \kappa_1 (q_\xi)^2 - \kappa_2 (q_\eta)^2 - 2\kappa_1 p_\xi q_\xi + 2\kappa_2 \partial_\xi^{-1}(p_\eta q_\xi)_\eta + v_0. \quad (4.35)$$

It follows from (4.32) that Eq. (4.13) for the invariant w_2 in terms of the variables p and q becomes

$$\begin{aligned} (p_{\xi\eta} + p_\eta q_\xi)_t &= \kappa_1 (p_{\xi\eta} + p_\eta q_\xi)_{\xi\xi} - \kappa_2 (p_{\xi\eta} + p_\eta q_\xi)_{\eta\eta} - \\ &- 2\kappa_1 ((p_{\xi\eta} + p_\eta q_\xi)(p_\xi - q_\xi))_\xi + 2\kappa_2 ((p_{\xi\eta} + p_\eta q_\xi)(p_\eta - q_\eta))_\eta. \end{aligned} \quad (4.36)$$

Equations (4.34)–(4.36) are compatible if the variable v_0 satisfies the relation

$$v_0 \xi_\eta + p_\eta v_0 \xi + q_\xi v_0 \eta = 0. \quad (4.37)$$

In the case where $v_0 \equiv 0$, system of three equations (4.34)–(4.36) implies the equivalent system of two equations

$$\begin{aligned} p_t &= \kappa_1 p_{\xi\xi} - \kappa_2 p_{\eta\eta} - \kappa_1 (p_\xi)^2 + \kappa_2 (p_\eta)^2 - 2\kappa_2 p_\eta q_\eta + 2\kappa_1 \partial_\eta^{-1} (p_\eta q_\xi)_\xi, \\ q_t &= -\kappa_1 q_{\xi\xi} + \kappa_2 q_{\eta\eta} + \kappa_1 (q_\xi)^2 - \kappa_2 (q_\eta)^2 - 2\kappa_1 p_\xi q_\xi + 2\kappa_2 \partial_\xi^{-1} (p_\eta q_\xi)_\eta. \end{aligned} \quad (4.38)$$

This system of equations was first obtained in a different context in [23].

To conclude this section, we discuss Miura-type transformations between different systems of nonlinear second-order DS-type equations obtained above in different gauges. For convenience, we let $P \equiv p$ and $Q \equiv q$ denote the solutions of system of nonlinear DS-type equations (4.26). We use the invariants w_1 and w_2 to obtain the relations between the variables P and Q of DS system (4.26) and the variables p and q of system (4.29), (4.30):

$$w_1 = (\log Q)_{\xi\eta} = p_{\xi\eta} - q_{\xi\eta}, \quad w_2 = PQ = -p_{\xi\eta}. \quad (4.39)$$

Relation (4.39) implies the corresponding Miura-type transformation

$$Q = e^{p-q}, \quad P = -p_{\xi\eta} e^{q-p}. \quad (4.40)$$

Quite similarly, for the pair of systems of DS-type equations (4.26) and (4.38), we obtain

$$w_1 = (\log Q)_{\xi\eta} = p_{\xi\eta} - q_{\xi\eta}, \quad w_2 = PQ = -p_{\xi\eta} - p_\eta q_\xi. \quad (4.41)$$

From (4.41), we also obtain a Miura-type transformation between the corresponding solutions

$$Q = e^{p-q}, \quad P = -(p_{\xi\eta} + p_\eta q_\xi) e^{q-p}. \quad (4.42)$$

Transformations (4.40) and (4.42) permit obtaining solutions of the well-known system of DS equations (4.26) from the corresponding solutions of systems of Eqs. (4.29), (4.30), and (4.38). Therefore, these transformations are Miura-type transformations between the solutions of gauge-equivalent systems of second-order DS-type equations.

5 Manifestly gauge-invariant system of integrable Nizhnik–Veselov–Novikov equations

We consider the results recently obtained in [17] concerning the gauge-invariant formulation of two-dimensional nonlinear evolution equations integrable using the two auxiliary linear problems

$$L_1 \psi = (\partial_{\xi\eta}^2 + u_1 \partial_\xi + v_1 \partial_\eta + u_0) \psi = 0, \quad (5.1)$$

$$L_2 \psi = (\partial_t + \kappa_1 \partial_\xi^3 + \kappa_2 \partial_\eta^3 + u_2 \partial_\xi^2 + v_2 \partial_\eta^2 + \tilde{u}_1 \partial_\xi + \tilde{v}_1 \partial_\eta + v_0) \psi = 0, \quad (5.2)$$

where κ_1 and κ_2 are constants, $\xi = x + \sigma y$ and $\eta = x - \sigma y$ are spatial coordinates, and $\sigma^2 = \pm 1$.

We use the compatibility condition in the Manakov triad form [20] (see (4.3)) to reduce auxiliary linear problems (5.1) and (5.2) to the previously obtained and well-known Nizhnik–Veselov–Novikov (NVN) equations and to some other equations [24–27] (also see [12, 13] and the references therein).

In the case under study, just as for the system of 2DgDLW equations, formulas (4.4) and (4.5) give the field variables, the classical invariants w_2 and w_1 , and the pure gauge variable ρ , which are convenient for the gauge-invariant formulation. In the case of auxiliary linear problem (5.2), the first invariant w_1 is equal to zero ($w_1 \equiv 0$), and the corresponding integrable system of nonlinear equations [17] in terms of the variables ρ and w_2 has the form

$$\rho_t = -\kappa_1 \rho_{\xi\xi\xi} - \kappa_2 \rho_{\eta\eta\eta} - 3\kappa_1 \rho_\xi \partial_\eta^{-1} w_{2\xi} - 3\kappa_2 \rho_\eta \partial_\xi^{-1} w_{2\eta} + v_0 \rho, \quad (5.3)$$

$$w_{2t} = -\kappa_1 w_{2\xi\xi\xi} - \kappa_2 w_{2\eta\eta\eta} - 3\kappa_1 (w_2 \partial_\eta^{-1} w_{2\xi})_\xi - 3\kappa_2 (w_2 \partial_\xi^{-1} w_{2\eta})_\eta. \quad (5.4)$$

System of equations (5.3), (5.4) in terms of the variables $\phi = \log \rho$ and w_2 becomes

$$\begin{aligned} \phi_t = & -\kappa_1 \phi_{\xi\xi\xi} - \kappa_2 \phi_{\eta\eta\eta} - \kappa_1 (\phi_\xi)^3 - \kappa_2 (\phi_\eta)^3 - 3\kappa_1 \phi_\xi \phi_{\xi\xi} - 3\kappa_2 \phi_\eta \phi_{\eta\eta} - \\ & - 3\kappa_1 \phi_\xi \partial_\eta^{-1} w_{2\xi} - 3\kappa_2 \phi_\eta \partial_\xi^{-1} w_{2\eta} + v_0, \end{aligned} \quad (5.5)$$

$$w_{2t} = -\kappa_1 w_{2\xi\xi\xi} - \kappa_2 w_{2\eta\eta\eta} - 3\kappa_1 (w_2 \partial_\eta^{-1} w_{2\xi})_\xi - 3\kappa_2 (w_2 \partial_\xi^{-1} w_{2\eta})_\eta. \quad (5.6)$$

Equation (5.4) (or (5.6)) for the gauge invariant $w_2 = u_0 - u_{1\xi} - u_1 v_1$ of the last two systems exactly coincides in form with the well-known NVN equation [24, 25]

$$u_t = -\kappa_1 u_{\xi\xi\xi} - \kappa_2 u_{\eta\eta\eta} - 3\kappa_1 (u \partial_\eta^{-1} u_\xi)_\xi - 3\kappa_2 (u \partial_\xi^{-1} u_\eta)_\eta. \quad (5.7)$$

Integrable system (5.3), (5.4) (or (5.5), (5.6)) can therefore be called the system of NVN equations.

Integrable system of NVN equations (5.3), (5.4) (or (5.5), (5.6)) has the following gauge structure:

- a. It contains manifestly gauge-invariant equation (5.4) (or (5.6)) for the invariant w_2 .
- b. It contains Eq. (5.3) (or (5.5)) for the pure gauge variable ρ (or ϕ) with some additional terms containing the gauge invariant w_2 and the additional field variable v_0 used in auxiliary linear problem (5.2).

For $w_2 = 0$, system of NVN equations (5.3), (5.4) (or (5.5), (5.6)) reduces to the single linear equation

$$\rho_t = -\kappa_1 \rho_{\xi\xi\xi} - \kappa_2 \rho_{\eta\eta\eta} + v_0 \rho. \quad (5.8)$$

In terms of the variable $\phi = \log \rho$, linear equation (5.8) looks like the third-order Burgers equation

$$\phi_t = -\kappa_1 \phi_{\xi\xi\xi} - \kappa_2 \phi_{\eta\eta\eta} - \kappa_1 (\phi_\xi)^3 - \kappa_2 (\phi_\eta)^3 - 3\kappa_1 \phi_\xi \phi_{\xi\xi} - 3\kappa_2 \phi_\eta \phi_{\eta\eta} + v_0, \quad (5.9)$$

which is linearized by the substitution $\phi = \log \rho$ and is hence C -integrable.

We let $C(\phi, u_0, v_0)$ denote the gauge corresponding to the nonzero field variables $u_1 = \phi_\eta$, $v_1 = \phi_\xi$, u_0 , and v_0 of auxiliary linear problems (5.1) and (5.2) and hence to system of NVN equations (5.5), (5.6) in general position. In different gauges, the NVN system implies

different gauge-equivalent integrable nonlinear equations. The solutions of these equations are related by Miura-type transformations.

For example, in the gauge $C(0, u_0, 0)$, system of NVN equations (5.5), (5.6) reduces to the well-known NVN equation [24, 25] for the field variable u_0 :

$$u_{0t} = -\kappa_1 u_{0\xi\xi\xi} - \kappa_2 u_{0\eta\eta\eta} - 3\kappa_1 (u_0 \partial_\eta^{-1} u_{0\xi})_\xi - 3\kappa_2 (u_0 \partial_\xi^{-1} u_{0\eta})_\eta. \quad (5.10)$$

In another gauge, $C(\phi, 0, v_0)$, NVN system (5.5), (5.6) becomes

$$\begin{aligned} \phi_t = & -\kappa_1 \phi_{\xi\xi\xi} - \kappa_2 \phi_{\eta\eta\eta} - \kappa_1 (\phi_\xi)^3 - \kappa_2 (\phi_\eta)^3 + \\ & + 3\kappa_1 \phi_\xi \partial_\eta^{-1} (\phi_\xi \phi_\eta)_\xi + 3\kappa_2 \phi_\eta \partial_\xi^{-1} (\phi_\xi \phi_\eta)_\eta + v_0, \end{aligned} \quad (5.11)$$

$$\begin{aligned} (\partial_{\xi\eta}^2 + \phi_\eta \partial_\xi + \phi_\xi \partial_\eta) \phi_t = & (\partial_{\xi\eta}^2 + \phi_\eta \partial_\xi + \phi_\xi \partial_\eta) \left[-\kappa_1 \phi_{\xi\xi\xi} - \kappa_2 \phi_{\eta\eta\eta} - \right. \\ & \left. - \kappa_1 (\phi_\xi)^3 - \kappa_2 (\phi_\eta)^3 + 3\kappa_1 \phi_\xi \partial_\eta^{-1} (\phi_\xi \phi_\eta)_\xi + 3\kappa_2 \phi_\eta \partial_\xi^{-1} (\phi_\xi \phi_\eta)_\eta \right]. \end{aligned} \quad (5.12)$$

By (5.11) and (5.12), NVN system (5.5), (5.6) in the gauge $C(\phi, 0, v_0)$ reduces to the system of equations

$$\begin{aligned} \phi_t = & -\kappa_1 \phi_{\xi\xi\xi} - \kappa_2 \phi_{\eta\eta\eta} - \kappa_1 (\phi_\xi)^3 - \kappa_2 (\phi_\eta)^3 + \\ & + 3\kappa_1 \phi_\xi \partial_\eta^{-1} (\phi_\xi \phi_\eta)_\xi + 3\kappa_2 \phi_\eta \partial_\xi^{-1} (\phi_\xi \phi_\eta)_\eta + v_0, \\ (\partial_{\xi\eta}^2 + \phi_\eta \partial_\xi + \phi_\xi \partial_\eta) v_0 = & 0. \end{aligned} \quad (5.13)$$

For $v_0 = 0$, system of equations (5.13) reduces to the well-known modified NVN (mNVN) equation

$$\phi_t = -\kappa_1 \phi_{\xi\xi\xi} - \kappa_2 \phi_{\eta\eta\eta} - \kappa_1 (\phi_\xi)^3 - \kappa_2 (\phi_\eta)^3 + 3\kappa_1 \phi_\xi \partial_\eta^{-1} (\phi_\xi \phi_\eta)_\xi + 3\kappa_2 \phi_\eta \partial_\xi^{-1} (\phi_\xi \phi_\eta)_\eta, \quad (5.14)$$

which was first obtained in a different context by Konopelchenko [26]. We note that mNVN equation (5.14) differs from the mNVN equation obtained by Bogdanov [27]. The new system of equations (5.13) can also be called a system of mNVN equations.

Obviously, the solutions u_0 and ϕ of NVN equations (5.10) and mNVN equations (5.14) are related by the Miura-type transformation

$$u_0 = -\phi_{\xi\eta} - \phi_\xi \phi_\eta \quad (5.15)$$

through the gauge invariant $w_2 = u_0 = -\phi_{\xi\eta} - \phi_\xi \phi_\eta$ (calculated in the different gauges $C(0, u_0, 0)$ and $C(\phi, 0, 0)$).

In the one-dimensional limit with coinciding derivatives $\partial_\xi = \partial_\eta$, mNVN equation (5.14) reduces to mKdV equation in potential form

$$\phi_t = -\kappa \phi_{\xi\xi\xi} + 2\kappa (\phi_\xi)^3, \quad (5.16)$$

where $\kappa = \kappa_1 + \kappa_2$. In terms of the variable $v_1 = \phi_\xi$, this is the mKdV equation

$$v_{1t} = -\kappa v_{1\xi\xi\xi} + 6\kappa v_1^2 v_{1\xi}. \quad (5.17)$$

6 Conclusion

The ideas of gauge invariance are currently widely used in the theory of integrable nonlinear equations. A manifestly gauge-invariant formulation of integrable systems of nonlinear equations can be given in all cases where the gauge freedom is properly taken into account in the corresponding auxiliary linear problems.

Acknowledgments. This work is supported by the Analytic Departmental Goal-Oriented Program of the Russian Federation Ministry of Science and Education “Development of the Higher School Potential (2009–2010)” (Grant No. 2.1.1/1958) and Novosibirsk State Technical University (Grant in Basic Research 2008–2009).

References

- [1] V. E. Zakharov and A. B. Shabat, *Funct. Anal. Appl.*, **8**, 226–235 (1974).
- [2] E. A. Kuznetsov and A. V. Mikhailov, *Theor. Math. Phys.*, **30**, 193–200 (1977).
- [3] V. E. Zakharov and A. V. Mikhailov, *Sov. Phys. JETP*, **47**, 1017–1027 (1978).
- [4] V. E. Zakharov and L. A. Takhtadzhyan, *Theor. Math. Phys.*, **38**, 17–23 (1979).
- [5] B. G. Konopelchenko, *Phys. Lett. A*, **92**, 323–327 (1982).
- [6] B. G. Konopelchenko and V. G. Dubrovsky, *Phys. Lett. A*, **95**, 457–462 (1983).
- [7] B. G. Konopelchenko and V. G. Dubrovsky, *Ann. Phys.*, **156**, 265–302 (1984).
- [8] V. E. Zakharov, S. V. Manakov, S. P. Novikov, and L. P. Pitaevskii, *Theory of Solitons: The Inverse Scattering Method* [in Russian], Nauka, Moscow (1980); English transl., Plenum, New York (1984).
- [9] L. A. Takhtadzhyan and L. D. Faddeev, *Hamiltonian Approach in the Theory of Solitons* [in Russian], Nauka, Moscow (1986); English transl.: L. D. Faddeev and L. A. Takhtadzhyan, *The Hamiltonian Methods in the Theory of Solitons*, Berlin, Springer (1987).
- [10] M. J. Ablowitz and P. A. Clarkson, *Solitons, Nonlinear Evolution Equations, and Inverse Scattering* (London Math. Soc. Lect. Note Ser., Vol. 149), Cambridge Univ. Press, Cambridge (1991).
- [11] B. G. Konopelchenko, *Nonlinear Integrable Equations: Recursion Operators, Group-Theoretical and Hamiltonian Structures of Soliton Equations* (Lect. Notes Phys., Vol. 270), Springer, Berlin (1987).
- [12] B. G. Konopelchenko, *Introduction to Multidimensional Integrable Equations: The Inverse Spectral Transform in 2+1 Dimensions*, Plenum, New York (1992).
- [13] B. G. Konopelchenko, *Solitons in Multidimensions: Inverse Spectral Transform Method*, World Scientific, Singapore (1993).

- [14] B. G. Konopelchenko and C. Rogers, “Bäcklund and reciprocal transformations: Gauge connections,” in: *Nonlinear Equations in the Applied Sciences* (Math. Sci. Engrg., Vol. 185, W. F. Ames and C. Rogers, eds.), Acad. Press, New York (1992), pp. 317–362.
- [15] V. S. Dryuma, *JETP Lett.*, **19**, 387–388 (1974).
- [16] B. G. Konopelchenko and V. G. Dubrovsky, *Phys. Lett. A*, **102**, 15–17 (1984).
- [17] V. G. Dubrovsky and A. V. Gramolin, *J. Phys. A*, **41**, 275208 (2008). Also available as arXiv:0802.2334.
- [18] M. Boiti, J. J. P. Leon and F. Pempinelli, *Inverse Problems*, **3**, 371–387 (1987).
- [19] A. Davey and K. Stewartson, *Proc. Roy. Soc. London Ser. A*, **338**, 101–110, (1974).
- [20] S. V. Manakov, *Russ. Math. Surveys*, **31**, 245–246 (1976).
- [21] A. R. Forsyth, *Theory of Differential Equations*, Vol. 6, Cambridge Univ. Press, Cambridge (1906).
- [22] L. J. F. Broer, *Appl. Sci. Res.*, **31**, 377–395 (1975).
- [23] B. G. Konopelchenko, *Inverse Problems*, **4**, 151–163 (1988).
- [24] L. P. Nizhnik, *Sov. Phys. Dokl.*, **25**, 706–708 (1980).
- [25] A. P. Veselov and S. P. Novikov, *Sov. Math. Dokl.*, **30**, 588–591 (1984).
- [26] B. G. Konopelchenko, *Rev. Math. Phys.*, **2**, 399–440 (1990).
- [27] L. V. Bogdanov, *Theor. Math. Phys.*, **70**, 219–223 (1987).